

# Effective interventions for mental health in male-dominated workplaces

Nicole K. Lee, Ann Roche, Vinita Duraisingam, Jane A. Fischer and Jacqui Cameron

Dr Nicole K. Lee is an Associate Professor, based at National Centre for Education and Training on Addiction (NCETA), Flinders University, Adelaide, Australia and National Drug Research Institute, Curtin University, Perth, Australia. Professor Ann Roche is the Director, Vinita Duraisingam is a Project Manager/Psychologist and Jane A. Fischer is a Researcher, all are based at National Centre for Education and Training on Addiction, Flinders University, Adelaide, Australia. Jacqui Cameron is a Researcher, based at National Centre for Education and Training on Addiction, Flinders University, Adelaide, Australia and Department of General Practice, University of Melbourne, Melbourne, Australia.

This project was part of a larger project funded by *beyondblue* and the Australian Government Department of Health. The full report from this project is also available: Roche, A.M., Fischer, J., Pidd, K., Lee, N.K., Battams, S. and Nicholas, R. (2012), "Workplace mental illness and substance use disorders in male-dominated industries: A Systematic Literature Review", National Centre for Education and Training on Addiction, Flinders University, Adelaide, [www.beyondblue.org.au](http://www.beyondblue.org.au)

## Abstract

**Purpose** – The purpose of this paper is to identify mental health interventions within male-dominated industries.

**Design/methodology/approach** – A systematic literature review was undertaken, examining mental health interventions within male-dominated industries. Major electronic databases, grey literature and reference lists for English language studies published January 1990-June 2012 were searched. Independent extraction of the studies was completed by two reviewers using predefined data fields including study quality measures.

**Findings** – Five studies met inclusion criteria. The available evidence suggests that effective interventions to address anxiety and depression in male-dominated industries include: improving mental health literacy and knowledge, increasing social support, improving access to treatment, providing education for managers and addressing workload issues.

**Practical implications** – Working conditions and the workplace can have a significant impact on a worker's mental health. Work-related factors including working conditions, job demands and social support in the workplace are particularly important for the mental health workers. Indeed, poor work conditions have been associated with poorer mental health outcomes in particular anxiety and depression, however, little work has been conducted on mental health interventions in the workplace and further the impact on male-dominated industries.

**Originality/value** – Overall, the body of evidence supporting effective interventions for mental health problems among workers in male-dominated industries is limited. Nonetheless, the evidence does suggest that mental health interventions in male-dominated industries is logistically feasible and can have some positive impact on the mental health of workers, particularly for high prevalence low severity disorders such as anxiety and depression.

**Keywords** Depression, Mental health, Anxiety, Male-dominated workplaces, Work related health

**Paper type** Literature review

## Introduction

The substantial impact that the workplace can have on a worker's mental health is now well established (Faragher *et al.*, 2005; Maslach, 2003; Segerstrom and Miller, 2004). The effect on mental health can be both positive, by providing regular activity, time structure, social contact, a sense of collective effort and social identity (Cercarelli *et al.*, 2012) and negative, as a source of psychological stress related to job demands and lack of social support in the workplace. Poor work conditions have been associated with even poorer mental health than among those who are unemployed (Butterworth *et al.*, 2011; Kuoppala *et al.*, 2008; Stansfeld and Candy, 2006).

The workplace bears a substantial proportion of the social and economic burden associated with mental illness (Conti and Burton, 1994; Dewa and Lin, 2000). Poor mental health is estimated to be associated with 50-60 per cent of all workplace absenteeism (Milczarek *et al.*, 2009) and the annual cost of mental health disorders is estimated to be around \$20 billion in Australia (Australian Bureau of Statistics, 2009; World Health Organization, 2003) and more than \$100 million in the USA (National Center for Health Statistics, 2007), including lost

productivity and workforce participation (Burton *et al.*, 1999; Stewart *et al.*, 2003). In addition, physical health problems are over-represented among those with mental health issues, further compounding the effects on the individual and the workplace (National Mental Health Consumer & Carer Forum, 2010). The enormous cost of mental health disorders creates a social and economic imperative to reduce the prevalence of mental health problems within the working population.

Workers affected by mental health problems are at substantially greater risk of becoming disengaged from the workforce, being less productive (Lim *et al.*, 2000), experiencing higher levels of physical health problems and being involved in accidents (Harnois and Gabriel, 2000).

Mental health problems are significant contributors to the global burden of disease more generally and comprise the largest contributor to years lost to disability, with depression (2.5 per cent) and anxiety (1.1 per cent) the main contributors (Murray *et al.*, 2013). From 1990 to 2010, major depressive disorders increased from 15th to 11th position (a 37 per cent increase) as a cause of Disability Adjusted Life Years (Murray *et al.*, 2013). Depression and anxiety are the most common mental health problems: globally, anxiety disorders have prevalence in the prior 12-month period of 2.4–18.2 per cent and mood disorders a 12-month prevalence of 0.8–9.6 per cent (Demyttenaere *et al.*, 2004).

In the general population, women have higher rates of anxiety and depression than men (22 per cent vs 18 per cent); however, a number of male-dominated industries (i.e. industries in which more than 70 per cent of workers are men (Australian Bureau of Statistics, 2008b) have higher than average rates of anxiety and mood disorders. These industries include agriculture (20.6 per cent), construction (23.3 per cent), mining (22.4 per cent) and utilities (20.7 per cent) (Australian Bureau of Statistics, 2008a).

Research on mental health problems in the workplace suggests that the personal and financial costs could be reduced if a greater proportion of workers who need treatment received it. However, men in particular are often reluctant to seek assistance for mental health concerns. They subsequently delay accessing treatment or help and attend mental health treatment at lower rates than women (Addis and Mahalik, 2003; Barney *et al.*, 2006; Galdas *et al.*, 2005), with 339 and 285 service contacts per 1,000 population for men and women, respectively in 2010–2011 in Australia.

Therefore, workers in male-dominated industries are at higher risk of mental health disorders but are less likely to access treatment than the general population (Battams *et al.*, 2014). The stigma that particularly men feel in addressing their mental health issues, especially in the workplace where jobs may be jeopardised, means that particular attention is needed on the types of interventions that are effective in these workplaces. The male-dominated workplace presents an excellent opportunity to access a difficult-to-reach population and to address mental health issues at work within a broader public health perspective (Lamontagne *et al.*, 2007; Pidd and Roche, 2008; Roche *et al.*, 2012).

However, despite a growing interest in, and attention on, men's mental health issues, little research has been undertaken to identify effective strategies for male-dominated workplaces. A systematic review was undertaken to examine the current evidence base for workplace interventions addressing mental health problems in male-dominated industries.

## Method

A systematic literature review was undertaken to identify and review studies examining mental health interventions within male-dominated industries. Due to the small number of papers and heterogeneity of measures to assess outcomes, a meta-analytic review was not possible; a narrative systematic review is therefore reported.

## Inclusion and exclusion criteria

Methods and results are reported in accordance with the PRISMA statement (Liberati *et al.*, 2009; Moher *et al.*, 2010). The analysis and inclusion criteria were specified in advance and

agreed upon by the research team using a predefined protocol (available from the lead author) that specified the process for data extraction (Von Elm *et al.*, 2007) as well as quality measures (National Center for Health Statistics, 2007) to be used to assess the quality of each study.

The definition of male-dominated industry was based on Australia Bureau of Statistics (Australian Bureau of Statistics, 2011) figures indicating the industries with the highest percentage of male workers (see Table I).

A male-dominated industry was determined using the Australian and New Zealand Standard Industrial Classification (ANZSIC) (Australian Bureau of Statistics, 2008b).

We defined a male-dominated industry as one in which there are greater than 70 per cent male workers. This definition was chosen to examine industries that are more-than-marginally male dominated, to ensure that the particular mental health issues for men, and of implementing interventions in male-dominated workplaces, were specifically examined. According to the ANZSIC, these industries are agriculture, construction, mining, manufacturing, transport and utilities. Although some other industries have a higher proportion of men than women, including public administration and safety, which includes fire, police and ambulance services, these industries had < 70 per cent men in the ANZSIC system (Australian Bureau of Statistics, 2006). Within this classification, military are not considered an industry *per se*, rather a workplace, and personnel are categorised by their industry type within this workplace.

Studies were included if they examined mental health interventions in the workplace in one of the defined male-dominated industries; were published between January 1990 and June 2012 in English; with a sample of male or female participants in paid formal work; and measured clinically significant anxiety or depression. Studies were excluded if they primarily investigated mental health issues other than anxiety and depression, including measures of subclinical symptoms only, or examined only volunteer workers or did not include workers in the pre-defined male-dominated industries.

### Search strategy

Searches were undertaken using the following electronic databases: CINAHL, Cochrane Database of Systematic Reviews, PubMed, PsycINFO, Informit and Scopus.

**Table I** B43 Industry of employment by sex

Industry (ABS, 2011)	Total	Males	
	<i>n</i>	<i>n</i>	%
Construction	828,910	719,206	86.8
Mining	176,563	145,765	82.6
Transport, postal and warehousing	479,181	368,050	76.8
Electricity, gas, water and waste services	115,611	87,954	76.1
Manufacturing	902,829	668,017	74.0
Agriculture, forestry and fishing	249,828	174,774	70.0
Wholesale trade	403,801	264,640	65.5
Information media and telecommunications	178,190	103,637	58.2
Professional, scientific and technical services	730,062	404,357	55.4
Public administration and safety	689,931	372,301	54.0
Arts and recreation services	151,574	79,358	52.4
Rental, hiring and real estate services	158,853	78,003	49.1
Administrative and support services	323,779	156,926	48.5
Financial and insurance services	377,352	176,626	46.8
Accommodation and food services	650,397	284,815	43.8
Retail trade	1,057,310	446,002	42.2
Education and training	804,418	240,824	29.9
Health care and social assistance	1,167,634	245,316	21.0
Other services	378,217	211,298	55.9
Inadequately described/Not stated	233,885	138,800	59.4
Total	10,058,325	5,366,669	53.4

An example of the string search used is provided below:

1. mental health.mp. or exp Mental Health/
2. mental health.m\_titl
3. 1 or 2
4. exp Anxiety Disorders/or exp Anxiety/or anxiety.mp
5. anxiety.m\_titl
6. 4 or 5
7. depression.mp. or exp Major Depression/
8. depression.m\_titl
9. 7 or 8
10. exp Major Depression/or exp Psychotherapy/or generalised anxiety.mp. or exp Risk Factors/or exp Anxiety/or exp Anxiety Disorders/or exp Generalized Anxiety Disorder/or exp Treatment Effectiveness Evaluation/or exp Panic Disorder/
11. generalised anxiety.m\_titl
12. 10 or 11
13. mental health/or well being/
14. "depression (emotion)"/
15. anxiety/or agitation/or anxiety disorders/or generalized anxiety disorder/or panic/or panic attack/or panic disorder/
16. 13 or 14 or 15
17. 3 or 6 or 9 or 12 or 16

A total of  $n = 282$  abstracts were retrieved and assessed for inclusion in this systematic review. Of these  $n = 238$  were further excluded as they did not satisfy the inclusion criteria. After data extraction and quality assessment a further  $n = 39$  studies were excluded leaving the final number of included studies ( $n = 5$ ). Of these, three were assessed to be of good methodological quality (Figure 1).


### ***Screening, data extraction and quality assessment***

Studies were screened twice. First, two reviewers screened the title and abstract of each article against inclusion criteria. At the second screen, one reviewer screened the full article and checked that the study met the specified inclusion criteria. This process was then checked and confirmed by a second reviewer. A data extraction codebook was developed to ensure consistency between reviewers and any disagreement for inclusion was referred to a third reviewer.

Study quality was assessed in two ways. The level of evidence of the study was considered against the National Health and Medical Research Council's evidence hierarchy (NHMRC, 1999, 2000).

To assess for risk of bias, the *Quality Assessment Tool for Quantitative Studies – Effective Public Health Practice Project* was used (National Collaborating Centre for Methods and Tools, 2009), which assesses selection bias, study design, confounders, blinding, data collection methods, withdrawals and drop-outs, intervention integrity and analyses. A global rating was derived for each study, ranging from “weak” to “strong”. Studies with no ratings of “weak” and at least four ratings of “strong” for any of the assessment criteria were considered methodologically strong. Studies that had one weak rating and less than four strong ratings were assessed as

**Figure 1** Flow of information through the different phases of a systematic review on mental health interventions in male-dominated industries



methodologically moderate. Studies that had two or more weak ratings were assessed as methodologically weak.

To ensure consistency in data extraction a data extraction template was developed based on the STROBE statement (Von Elm *et al.*, 2007). It included methods (selection of subjects, allocation and blinding, confounders and statistical analyses), results, conflict of interest and other sources of bias.

## Results

Five studies met the inclusion criteria and these are summarised in Table II.

Most studies used multi-modal interventions, making it difficult to determine which components of the intervention were most effective. The quality of the studies varied widely and did not measure clinical mental health symptoms, focusing instead on health and wellbeing as outcome measures. Three of the five studies were assessed to be of good methodological quality. Of the five included studies, three were undertaken in Japan and two in Finland, potentially limiting the generalisability of the work.

A group of  $n = 782$  manufacturing workers who scored  $> 8$  on the General Health Questionnaire (GHQ-30) in Japan were followed up every six months for one year, without any specific intervention, and the relationship between job stress and recovery from mental illness was examined (Mino *et al.*, 2000). Of the 232 workers at the six-month follow-up, 113 (49 per cent)

**Table II** Included mental health intervention studies

Study	n	Participants	Intervention	Outcome measures	Follow-up	Outcomes	Quality assessment
Mino <i>et al.</i> (2000), Japan Cohort study	n = 782	Mean Age Not reported Gender (%male) 49%	No formal intervention but examined the relationship between job stress and indicators on recovery or improvement from mental ill health	General health questionnaire Perceived job stress Family life satisfaction Physical health	6 months	Outcome 1: Recovery proportion at 6 months: 49% (n = 113). Recovery proportion at 1 year: 66% (n = 36). No significant associations between recovery and gender, age, family life satisfaction or physical health Outcome 2: No significant associations between recovery and job stress during first 6 months. Significant associations found between recovery and having too much responsibility – OR 4.16 (1.3-13.1) (95% CI, p < 0.05) Authors' conclusion: Relief from excessive responsibility at work may enhance recovery in mentally ill workers Outcome 1: Sickness absence day rates due to mental disorders had a significant reduction from 0.06 to 0.03% (p < 0.001) For short sickness absence, day rates due to mental disorders increased from 0.01 to 0.03% but for long sickness absence, the day rate decreased from 0.05 to 0%	Fair
Shimizu <i>et al.</i> (2003), Japan Case study	n = 1,259	Mean Age 37.6 ± 9.6 Gender (%male) 100%	Mental health care program (MCAp) aimed to reduce absence due to mental health disorders and included three elements – social support, treatment support, and managerial education	Sickness absence days	2.5 years	Authors' conclusion: In conclusion, the trends in sickness absence due to mental health disorders change in the 2.5 years after the introduction of a mental health care program Outcome 1: Mean differences in sickness absence was 11 days (95% CI, 1-20) The intervention was effective in reducing sickness absence Outcome 2:	Poor
Taimela <i>et al.</i> (2008), Finland RCT	n = 1,341	Mean Age Not reported Gender (%male) 88%	Intervention participants received a targeted occupational health intervention program, which included consultation and personalised feedback from occupational	Sickness absence days Sleep disturbances Work-related stress Fatigue Depression Pain	1 year		Good

(continued)

**Table II**

Study	n	Participants	Intervention	Outcome measures	Follow-up	Outcomes	Quality assessment
Taimela <i>et al.</i> (2010), Finland RCT	n = 382	Mean Age 49 ± 9.4 Gender (%male) 93%	medical staff Participants could also attend for a consultation to a local occupational health service. Control participants could consult medical staff on request but received no feedback	Disability Working ability		After 12 months, the mean of sickness absence was 30 days in the control group (n = 192) and 11 days less (95% CI 1 to 20 days) in the intervention group (n = 192). No significant differences in self-reported work ability, depression, fatigue, stress, pain and physical impairment between groups, however depression and fatigue tended to be more present in the control group at 12 months Authors' conclusion: Workers at a high risk for sickness absence can be identified. Appropriate occupational health intervention can be effective in reducing sickness absence among high-risk workers Outcome 1: No modifying effect of age, gender, work status, depressive symptoms or previous sickness absence was found in the previous 12 months Outcome 2: The intervention was more effective for workers who reported comorbidities, i.e. more than one health problem (–22.5 days; 95% CI –35.5 to –9.5) and those who perceived an inability to work in the future due to health-related reasons (–7.4 days; 95% CI –10.5 to –4.3) Authors' conclusion: Occupational interventions can have a greater effect for certain workers such as those who have co-morbidities and those who perceive that they may be unable to work in the future due to health impairment	Good
Tsutsumi <i>et al.</i> (2009), Japan RCT	n = 97	Intervention Mean Age 49 Gender (%male) 51% Control Mean Age 41 Gender (%male) 69%	Participatory workplace mental health and productivity program (problem-solving, team-based approach on improving work environment and job processes)	General Health WHO Health and Work Performance Age	1 year	Outcome 1: GHQ scores remained unchanged for the intervention group but increased for the control group (t = –2.43, p = 0.020) GHQ scores at 12 months decreased for intervention (t = –2.37; p = 0.022) and increased for control group (t = 2.18; p = 0.035) with a significant interaction effect Outcome 2: HPQ scores increased in the intervention group (t = –2.37, p = 0.348) but decreased in the control group (t = 2.13, p = 0.040) Authors' conclusion: The participatory approach may be effective in improving mental health outcomes in workers	Good

had a score of  $<8$  on the GHQ-30 suggesting some recovery, the recovery rate reported at 12 months was 66 per cent. A relationship between job responsibility and recovery was identified and it was concluded that relief from excessive responsibility at work might enhance recovery in mentally ill workers.

A case study of an organisational mental health care programme (MCaP) was examined years before and after the introduction of a two and a half year mental health care programme, which included social support, treatment support and managerial education on mental health issues. The sample included  $n=1,259$  male workers in an industrial chemical manufacturing plant, outcome measures included sickness absences, duration of sickness absence and the sickness absence day rate. Mental health was identified using the ICD-10. The results identified significant reductions in sickness absence days (0.06-0.03 per cent) – shorter absences increased slightly but longer absences decreased significantly (0.05-0 per cent) (Shimizu *et al.*, 2003).

A study in Finland randomly assigned  $n=1,341$  (49 per cent construction industry) workers (88 per cent male) with high and intermediate risk of sickness absence to a targeted occupational health programme or usual care (Taimela *et al.*, 2008). Intervention participants received an 11-month targeted occupational health intervention programme, which included consultation and personalised feedback from occupational medical staff. Control participants could consult medical staff on request but received no feedback. There was a significant reduction in sickness absence in the high risk but not the intermediate risk group. No significant differences in self-reported work ability, depression, fatigue, stress, pain and physical impairment between groups, however depression and fatigue tended to be more present in the control group at 12 months. The authors concluded that identifying and responding to people at high risk of sickness absence is effective in reducing absences.

The same research team using a subsample ( $n=382$ ) of the same data as Taimela *et al.* (2008) and completed further analyses which found a significant interaction in the intervention group on self-rated future working ability, severity of physical impairment and co-morbidity. The other modifying effects from age, gender, occupational status, depression score and sickness absence in previous 12 months were not statistically significant. They concluded that the intervention was especially effective for workers who were certain that they would not be able to continue working in their current job due to health-related reasons, multiple health problems or severe physical impairment at work (Taimela *et al.*, 2010).

In one randomised control study ( $n=97$ ) 11 assembly lines (six intervention and five control) were randomly assigned to a participatory workplace mental health and productivity programme for 12 months, which included problem solving and team based approaches to improving work environments and job processes, or to a no intervention control group and followed up 12 months later (Tsutsumi *et al.*, 2009). Outcome measures included self-report on mental health measured by the GHQ and work performance measured by the WHO Health and Work Performance Questionnaire (HPQ). GHQ scores remained the same for the intervention group, whereas they increased for the control group but were not significant, the HPQ scores increased for the intervention group but decreased in the control group and this change was significant, at follow up the GHQ scores reduced and the intervention group increased. The authors suggest that this programme had a protective effect on mental health and job performance.

## Discussion

Overall, the body of evidence supporting effective interventions for mental health problems among workers in male-dominated industries is limited. Nonetheless, the evidence does suggest that mental health interventions in male-dominated industries are logistically feasible and can have some positive impact on the mental health of workers, particularly for high prevalence low severity disorders such as anxiety and depression.

The available evidence indicates that effective strategies to address anxiety and depression among workers in male-dominated industries include:

- the distribution of information to workers about mental health issues;
- providing additional social support;



- offering access to treatment and advice for workers;
- education for managers about mental health in the workplace; and
- specifically targeting intervention at groups at high risk for absenteeism. Addressing excessive workloads and providing relief periods from heavy workloads.

This systematic review is the first of its kind that has attempted to identify the available evidence for interventions to address mental health problems among workers in male-dominated industries. Five studies met the inclusion criteria. The studies were variable in their design and interventions, but some were of good quality.

Few studies have identified intervention strategies specifically for male-dominated industries but a number of studies have examined interventions for mental health in the workplace more generally (Barry *et al.*, 2009; Cooper and Cartwright, 1994; Giga *et al.*, 2003; Lamontagne *et al.*, 2007). The workplace holds considerable promise as a setting to introduce strategies that will prevent and/or ameliorate mental health problems among a largely difficult-to-access population. Opportunities for primary and secondary prevention of mental health problems are of paramount importance (Barry *et al.*, 2009).

Historically, mental-health-related workplace interventions have targeted the individual worker with varying degrees of success and inconclusive long-term outcomes (Cooper and Cartwright, 1994, 1997; van der Klink *et al.*, 2003). Alternatively, primary prevention through “proactive”, organisation-directed activities (e.g. increased social support and job control in the workplace), to circumvent the need for “reactive” secondary (e.g. stress management) and tertiary (e.g. employee assistance programmes) prevention has been suggested (Cooper and Cartwright, 1994).

Interventions that targeted the whole workplace, utilised team-based approaches, and used multiple strategies appeared to be effective. Most studies used a whole of workplace approach and implemented a menu of interventions, targeting individuals, teams and systems. This is consistent with general mental health workplace research that suggests that, although individually focused approaches can favourably affect individual-level outcomes, they do not influence organisational level change (Lamontagne *et al.*, 2007) overall; while organisational approaches have benefits at both individual and organisational levels, and offer greater scope for the prevention of mental health problems in male-dominated industries (Cooper and Cartwright, 1994). However, it remains difficult to determine which individual components in the suite of intervention activities were required to achieve positive outcomes.

While most workers have relatively little control over workplace factors, at the organisational level there is considerable untapped latitude to address crucial factors that impact on the workers’ mental health and wellbeing.

The results of this review suggest that there is a potential benefit in providing primary intervention and secondary prevention activities for mental health disorders within the workplace setting. The type of research that is possible within a real-world workplace setting is generally broad, often tailored to the workplace and not well controlled, making generalisations to other work places difficult.

A systematic review of risk factors for mental health disorders (Battams *et al.*, 2014) in male-dominated industries noted that social support and team environment, job demand, job variety and control, workplaces can have a positive primary and secondary preventive influence on the mental health and wellbeing of employees.

More generally, research suggests that it is the workplace environment (Michie and Williams, 2003; Stansfeld and Candy, 2006), including job control (Bond and Bunce, 2001), and structural factors such as economic climate, labour market conditions and employment policies, job security (Marmot *et al.*, 2009; Meltzer *et al.*, 2010) that make the greatest contribution to worker wellbeing. By addressing these risk factors through well-designed interventions workplaces can have an impact on workers’ mental health, wellbeing and hence on productivity.

Growing emphasis has been placed on the duty of care of employers towards their employees through occupational health and safety (OHS) law and policy, including the duty to provide a safe workplace to promote both physical and psychological health. OHS has traditionally focused on

physical safety. There is increasing recognition of the need for a “psychologically safe” work environment (Dollard and Bakker, 2010).

The interventions studied to date that indicate promise are relatively inexpensive to design, implement and evaluate. The potential return on investment, and to the community at large, from such interventions is substantial.

As costs for mental health treatment (National Institute of Mental Health, 2013), life expectancy and retirement age continue to increase, and greater workforce participation extends into older age (Shannon, 2013), the workplace will become an increasingly important early intervention avenue to address public health issues such as high prevalence mental health issues. Ensuring that workers’ wellbeing is fostered is of greater imperative than ever.

A considerably improved evidence base would be helpful in informing the judicious selection of effective and appropriate interventions that may be applied in the workplace. Given the high prevalence of mental health problems in male-dominated industries, further research in these areas is identified as a pressing priority.

### Limitations

There were a limited number of studies found and the samples varied, particularly in relation to culture/country, gender and occupational categories. A range of measures were also used to assess anxiety and depression; this level of heterogeneity precluded a meta-analysis.

### Conclusions

Mental health problems have numerous sequelae, including an association with absenteeism, lost productivity and physical health issues; those working in male-dominated industries have higher than average rates of mental health issues. Men continue to present for treatment at lower rates than women, limiting access to opportunistic treatment by health professionals. This review suggests that workplaces can influence mental health and wellbeing and provide some support for the introduction of mental health interventions in the workplace. This is especially important in workplaces with populations that have low treatment seeking rates, such as men. However, to date, the evidence is limited and further research is needed to identify both programmes of interventions that are effective and to isolate specific active components of such programmes.

### References

- Addis, M.E. and Mahalik, J.R. (2003), “Men, masculinity, and the contexts of help seeking”, *American Psychologist*, Vol. 58 No. 1, pp. 5-14.
- Australian Bureau of Statistics (2006), *1292.0 – Australian and New Zealand Standard Industrial Classification (ANZSIC), 2006*, Commonwealth of Australia and Crown Copyright New Zealand, Canberra, available at: [www.abs.gov.au/ausstats/abs@.nsf/mf/1292.0%3E](http://www.abs.gov.au/ausstats/abs@.nsf/mf/1292.0%3E) (accessed 12 August 2013).
- Australian Bureau of Statistics (2008a), *4326.0 – National Survey of Mental Health and Wellbeing: Summary of Results, 2007*, Australian Bureau of Statistics, Sydney, available at: [www.abs.gov.au/ausstats/abs@.nsf/mf/4326.0%3E](http://www.abs.gov.au/ausstats/abs@.nsf/mf/4326.0%3E) (accessed 12 August 2013).
- Australian Bureau of Statistics (2008b), *Australian and New Zealand Standard Industrial Classification*, Commonwealth of Australia and Crown Copyright New Zealand, Canberra.
- Australian Bureau of Statistics (2009), *4102.0 – Australian Social Trends, March 2009*, 4102.0, Australian Bureau of Statistics, Canberra, available at: [www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4102.0Main+Features30March2009%3E](http://www.abs.gov.au/AUSSTATS/abs@.nsf/Lookup/4102.0Main+Features30March2009%3E) (accessed 9 October 2014).
- Australian Bureau of Statistics (2011), *B43 Industry of Employment by Age and Sex*, Australian Bureau of Statistics, Canberra, available at: [http://stat.abs.gov.au/Index.aspx?DataSetCode=ABS\\_CENSUS2011\\_B43%3E](http://stat.abs.gov.au/Index.aspx?DataSetCode=ABS_CENSUS2011_B43%3E) (accessed 9 October 2014).
- Barney, L.J., Griffiths, K.M., Jorm, A.F. and Christensen, H. (2006), “Stigma about depression and its impact on help-seeking intentions”, *Australian and New Zealand Journal of Psychiatry*, Vol. 40 No. 1, pp. 51-4.

- Barry, M., Canavan, R., Clarke, A., Dempsey, C. and O'Sullivan, M. (2009), *Review of Evidence-Based Mental Health Promotion and Primary/Secondary Prevention*, Department of Health, Galway.
- Battams, S., Roche, A.M., Fischer, J., Lee, N.K., Cameron, J. and Kostadinov, V. (2014), "Workplace risk factors for anxiety and depression in male-dominated industries: a systematic review", *Health Psychology and Behavioral Medicine*, Vol. 2 No. 1, pp. 983-1008.
- Bond, F.W. and Bunce, D. (2001), "Job control mediates change in a work reorganization intervention for stress reduction", *Journal of Occupational Health Psychology*, Vol. 6 No. 4, pp. 290-302.
- Burton, W.N., Conti, D.J., Chen, C.Y., Schultz, A.B. and Edington, D.W. (1999), "The role of health risk factors and disease on worker productivity", *Journal of Occupational and Environmental Medicine*, Vol. 41 No. 10, pp. 863-77.
- Butterworth, P., Leach, L.S., Strazdins, L., Olesen, S.C., Rodgers, B. and Broom, D.H. (2011), "The psychosocial quality of work determines whether employment has benefits for mental health: results from a longitudinal national household panel survey", *Occupational and Environmental Medicine*, Vol. 68 No. 11, pp. 806-12.
- Cercarelli, R., Allsop, S., Evans, M. and Velander, F. (2012), *Reducing Alcohol-Related Harm in the Workplace: An Evidence Review: Full Report*, Victorian Health Promotion Foundation, Melbourne.
- Conti, D.J. and Burton, W.N. (1994), "The economic impact of depression in a workplace", *Journal of Occupational Medicine*, Vol. 36 No. 9, pp. 983-8.
- Cooper, C.L. and Cartwright, S. (1994), "Healthy mind; healthy organization – a proactive approach to occupational stress", *Human Relations*, Vol. 47 No. 4, pp. 455-71.
- Cooper, C.L. and Cartwright, S. (1997), "An intervention strategy for workplace stress", *Journal of Psychosomatic Research*, Vol. 43 No. 1, pp. 7-16.
- Demyttenaere, K., Bruffaerts, R., Posada-Villa, J., Gasquet, I., Kovess, V., Lepine, J.P., Angermeyer, M.C., Bernert, S., de Girolamo, G., Morosini, P., Polidori, G., Kikkawa, T., Kawakami, N., Ono, Y., Takeshima, T., Uda, H., Karam, E.G., Fayyad, J.A., Karam, A.N., Mneimneh, Z.N., Medina-Mora, M.E., Borges, G., Lara, C., de Graaf, R., Ormel, J., Gureje, O., Shen, Y., Huang, Y., Zhang, M., Alonso, J., Haro, J.M., Vilagut, G., Bromet, E.J., Gluzman, S., Webb, C., Kessler, R.C., Merikangas, K.R., Anthony, J.C., Von Korff, M.R., Wang, P.S., Brugha, T.S., Aguilar-Gaxiola, S. and Lee, S. (2004), "Prevalence, severity, and unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys", *JAMA*, Vol. 291 No. 21, pp. 2581-90.
- Dewa, C.S. and Lin, E. (2000), "Chronic physical illness, psychiatric disorder and disability in the workplace", *Social Science and Medicine*, Vol. 51 No. 1, pp. 41-50.
- Dollard, M.F. and Bakker, A.B. (2010), "Psychosocial safety climate as a precursor to conducive work environments, psychological health problems, and employee engagement", *Journal of Occupational and Organizational Psychology*, Vol. 83 No. 3, pp. 579-99.
- Faragher, E.B., Cass, M. and Cooper, C.L. (2005), "The relationship between job satisfaction and health: a meta-analysis", *Occupational and Environmental Medicine*, Vol. 62 No. 2, pp. 105-12.
- Galdas, P.M., Cheater, F. and Marshall, P. (2005), "Men and health help-seeking behaviour: literature review", *Journal of Advanced Nursing*, Vol. 49 No. 6, pp. 616-23.
- Giga, S.I., Noblet, A.J., Faragher, B. and Cooper, C.L. (2003), "The UK perspective: a review of research on organisational stress management interventions", *Australian Psychologist*, Vol. 38 No. 2, pp. 158-64.
- Harnois, G. and Gabriel, P. (2000), *Mental Health and Work: Impact, Issues and Good Practice*, Mental Health Policy and Service Development Department of Mental Health and Substance Dependence Non Communicable Diseases and Mental Health, Geneva.
- Kuoppala, J., Lamminpää, A. and Husman, P. (2008), "Work health promotion, job well-being, and sickness absences-a systematic review and meta-analysis", *Journal of Occupational and Environmental Medicine*, Vol. 50 No. 11, pp. 1216-27.
- Lamontagne, A.D., Keegel, T., Louie, A.M., Ostry, A. and Landsbergis, P.A. (2007), "A systematic review of the job-stress intervention evaluation literature, 1990-2005", *International Journal of Occupational and Environmental Health*, Vol. 13 No. 3, pp. 268-80.
- Liberati, A., Altman, D.G., Tetzlaff, J., Mulrow, C., Gøtzsche, P.C., Ioannidis, J.P.A., Clarke, M., Devereaux, P.J., Kleijnen, J. and Moher, D. (2009), "The PRISMA statement for reporting systematic reviews and meta-analyses

of studies that evaluate health care interventions: explanation and elaboration", *Annals of Internal Medicine*, Vol. 151 No. 4, pp. W-65-W-94.

Lim, D., Sanderson, K. and Andrews, G. (2000), "Lost productivity among full-time workers with mental disorders", *The Journal of Mental Health Policy and Economics*, Vol. 3 No. 3, pp. 139-46.

Marmot, M., Siegrist, J., Theorell, T. and Feeney, A. (2009), "Health and the psychosocial environment at work", in Marmot, M. and Wilkinson, R. (Eds), *Social Determinants of Health – The Solid Facts*, 2nd ed., Oxford University Press, Oxford, pp. 105-31.

Maslach, C. (2003), "Job burnout new directions in research and intervention", *Current Directions in Psychological Science*, Vol. 12 No. 5, pp. 189-92.

Meltzer, H., Bebbington, P., Brugha, T., Jenkins, R., McManus, S. and Stansfeld, S. (2010), "Job insecurity, socio-economic circumstances and depression", *Psychological Medicine*, Vol. 40 No. 8, pp. 1401-7.

Michie, S. and Williams, S. (2003), "Reducing work related psychological ill health and sickness absence: a systematic literature review", *Occupational and Environmental Medicine*, Vol. 60 No. 1, pp. 3-9.

Milczarek, M., Schneider, E. and González, R.E. (2009), *OHS in Figures: Stress at Work-Facts and Figures. Luxembourg: European Agency for Safety and Health at Work*, European Agency for Safety and Health at Work, Luxembourg.

Mino, Y., Shigemi, J., Tsuda, T., Yasuda, N., Babazono, A. and Bebbington, P. (2000), "Recovery from mental ill health in an occupational setting: a cohort study in Japan", *Journal of Occupational Health*, Vol. 42 No. 2, pp. 67-71.

Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G. and Group, P. (2010), "Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement", *International Journal of Surgery*, Vol. 8 No. 5, pp. 336-41.

Murray, C.J.L., Vos, T., Lozano, R., Naghavi, M., Flaxman, A.D., Michaud, C., Ezzati, M., Shibuya, K., Salomon, J.A. and Abdalla, S. (2013), "Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010", *The Lancet*, Vol. 380 No. 9859, pp. 2197-223.

National Center for Health Statistics (2007), *Health, United States, 2007, with Chartbook on Trends in the Health of Americans*, National Center for Health Statistics, Hyattsville, MD.

National Collaborating Centre for Methods and Tools (2009), *Quality Assessment Tool for Quantitative Studies*, McMaster University, Hamilton.

National Institute of Mental Health (2013), *Mental Healthcare Costs for All Americans (1996-2006)*, National Institute of Mental Health, Bethesda, MD, available at: [www.nimh.nih.gov/Statistics/4MH\\_AM9603.shtml%3E](http://www.nimh.nih.gov/Statistics/4MH_AM9603.shtml%3E) (accessed 19 November 2013).

National Mental Health Consumer & Carer Forum (2010), *Issue: Physical Health Impacts of Mental Illness*, National Mental Health Consumer & Carer Forum, Australia.

NHMRC (1999), *A Guide to the Development, Implementation and Evaluation of Clinical Practice Guidelines*, National Health and Medical Research Council, Canberra, available at: [www.nhmrc.gov.au/guidelines/publications/cp30%3E](http://www.nhmrc.gov.au/guidelines/publications/cp30%3E) (accessed 9 October 2014).

NHMRC (2000), *How to Use the Evidence: Assessment and Application of Scientific Evidence*, National Health and Medical Research Council, Canberra.

Pidd, K. and Roche, A.M. (2008), "Changing workplace culture: an integrated model for the prevention and treatment of alcohol-related problems", in Moore, D. and Dietze, P. (Eds), *Drugs and Public Health: Australian Perspective on Policy and Practice*, Oxford University Press, Melbourne, pp. 49-59.

Roche, A.M., Fischer, J., Pidd, K., Lee, N.K., Battams, S. and Nicholas, R. (2012), *Workplace Mental Illness and Substance Use Disorders in Male-Dominated Industries: A Systematic Literature Review*, National Centre for Education and Training on Addiction, Flinders University, Adelaide.

Segerstrom, S.C. and Miller, G.E. (2004), "Psychological stress and the human immune system: a meta-analytic study of 30 years of inquiry", *Psychological Bulletin*, Vol. 130 No. 4, pp. 601-30.

Shannon, H.S. (2013), "Epidemiological studies of older workers: research questions and", in Venables, K. (Ed.), *Current Topics in Occupational Epidemiology*, Oxford University Press, Oxford, pp. 91-9.

- Shimizu, T., Nagashima, S., Mizoue, T. and Nagata, S. (2003), "A mental health care program and sickness absence in a Japanese manufacturing plant", *J Occup Health*, Vol. 45 No. 4, pp. 234-7.
- Stansfeld, S. and Candy, B. (2006), "Psychosocial work environment and mental health – a meta-analytic review", *Scandinavian Journal of Work, Environment and Health*, Vol. 32 No. 6, pp. 443-62.
- Stewart, W.F., Ricci, J.A., Chee, E., Hahn, S.R. and Morganstein, D. (2003), "Cost of lost productive work time among US workers with depression", *JAMA*, Vol. 289 No. 23, pp. 3135-44.
- Taimela, S., Aronen, P., Malmivaara, A., Sintonen, H., Tiekso, J. and Aro, T.E. (2010), "Effectiveness of a targeted occupational health intervention in workers with high risk of sickness absence: baseline characteristics and adherence as effect modifying factors in a randomized controlled trial", *Journal of Occupational Rehabilitation*, Vol. 20 No. 1, pp. 14-20.
- Taimela, S., Justen, S., Aronen, P., Sintonen, H., Laara, E., Malmivaara, A., Tiekso, J. and Aro, T. (2008), "An occupational health intervention programme for workers at high risk for sickness absence. Cost effectiveness analysis based on a randomised controlled trial", *Occupational and Environmental Medicine*, Vol. 65 No. 4, pp. 242-8.
- Tsutsumi, A., Nagami, M., Yoshikawa, T., Kog, I.K. and Kawakami, N.E. (2009), "Participatory intervention for workplace improvements on mental health and job performance among blue-collar workers: a cluster randomized controlled trial", *Journal of Occupational and Environmental Medicine*, Vol. 51 No. 5, pp. 554-63.
- van der Klink, J., Blonk, R., Schene, A.H. and Van Dijk, F.J.H. (2003), "Reducing long term sickness absence by an activating intervention in adjustment disorder: a cluster randomised control trial", *Journal of Occupational and Environmental Medicine*, Vol. 60 No. 6, pp. 429-37.
- Von Elm, E., Altman, D.G., Egger, M., Pocock, S.J., Gøtzsche, P.C. and Vandenbroucke, J.P. (2007), "The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies", *Preventive Medicine*, Vol. 45 No. 4, pp. 247-51.
- World Health Organization (2003), *Investing in Mental Health*, Department of Mental Health and Substance Dependence, World Health Organization, Geneva.

## About the authors

Dr Nicole K. Lee is an Associate Professor at the National Centre for Education and Training on Addiction at the Flinders University. She is a Practising Psychologist and also holds an adjunct appointment at the National Drug Research Institute, Curtin University. She holds postgraduate qualifications in education and a PhD in Psychology. She has worked in the mental health area for more than 24 years and has held positions as clinician, trainer, academic and service manager. She has extensive experience in undertaking systematic reviews and other clinical and translational research and has worked with the World Health Organisation and the United National Office of Drugs and Crime. She has published more than 100 journal articles, books, chapters and clinical guidelines. She has served on the board of the Australian Association for Cognitive and Behaviour Therapy (AACBT), the Alcohol and other Drugs Council of Australia (ADCA) and the Australasian Professional Society on Alcohol and other Drugs (APSAD). Dr Nicole K. Lee is the corresponding author and can be contacted at: nicole.lee@flinders.edu.au

Professor Ann Roche is the Director of the National Centre for Education and Training on Addiction (NCETA), Flinders University. Ann has over 30 years' experience in public health and has worked as a Researcher, Educator and Policy Analyst and has held academic posts at the University of Sydney, the University of Newcastle, the University of Queensland and the Flinders University. For the past 25 years, Ann's professional activities have focused largely on AOD issues with a specific interest in workforce development, professional practice change, best practice and research dissemination. She holds graduate level qualifications in education and a PhD in public health. Ann has published extensively in AOD-related public health areas, including over 100 papers and reports, with several books and book chapters. In addition, Ann has worked as a consultant to the World Health Organization, undertaken numerous consultancies for government and non-government bodies and has acted as an Adviser on a

wide range of committees in the AOD field. She is a past President of the Australasian Professional Society on Alcohol and other Drugs (APSAD).

Vinita Duraisingam is a Project Manager/Psychologist at the National Centre for Education and Training, Flinders University. She holds a Master of Organisational Psychology and is a Registered Psychologist. She has a background in organisational psychology and business management, has extensive project management skills and a strong track record in data analysis and report writing.

Jane A. Fischer is a Senior Research Officer at the National Centre for Education and Training, Flinders University. She holds a Bachelor of Arts (Anthropology) and a Master of Social Planning and Development Degree, and is currently completing a Doctorate of Philosophy in the quality of life of substance users. Jane has held public health academic positions in the alcohol, tobacco and other drugs field for 15 years in both university and government settings.

Jacqui Cameron holds a Master of Philosophy (MPhil) in Social Science Research and is a qualified Social Worker with demonstrated practice experience. She holds an adjunct appointment at the National Centre for Education and Training on Addiction (NCETA), Flinders University and Honorary Fellow appointment with the Department of General Practice, University of Melbourne. Cameron is a mixed method researcher with over 12 years of International research experience in social research and clinical evaluations. She also has experience in training and workshop facilitation in clinical settings.

---

To purchase reprints of this article please e-mail: [reprints@emeraldinsight.com](mailto:reprints@emeraldinsight.com)  
Or visit our web site for further details: [www.emeraldinsight.com/reprints](http://www.emeraldinsight.com/reprints)