

RESEARCH ARTICLE

The Potential for Mindfulness-Based Intervention in Workplace Mental Health Promotion: Results of a Randomized Controlled Trial

Shu-Ling Huang^{1,2}, Ren-Hau Li¹, Feng-Ying Huang³, Feng-Cheng Tang^{4,5*}

1 Department of Psychology, Chung-Shan Medical University, Taichung, Taiwan, **2** Room of Clinical Psychology, Chung-Shan Medical University Hospital, Taichung, Taiwan, **3** Department of Education, National Taipei University of Education, Taipei, Taiwan, **4** Department of Occupational Medicine, Changhua Christian Hospital, Changhua, Taiwan, **5** Department of Leisure Services Management, Chaoyang University of Technology, Taichung, Taiwan

* 106159@cch.org.tw



OPEN ACCESS

Citation: Huang S-L, Li R-H, Huang F-Y, Tang F-C (2015) The Potential for Mindfulness-Based Intervention in Workplace Mental Health Promotion: Results of a Randomized Controlled Trial. PLoS ONE 10(9): e0138089. doi:10.1371/journal.pone.0138089

Editor: James Coyne, University of Pennsylvania, UNITED STATES

Received: October 6, 2014

Accepted: August 24, 2015

Published: September 14, 2015

Copyright: © 2015 Huang et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was funded by the Health Promotion Administration, Ministry of Health and Welfare, Taiwan (HPA 100-Soc-4). (<http://www.hpa.gov.tw/BHPNet/Web/Index/Index.aspx>.) The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Objectives

This study aims to intensively evaluate the effectiveness of mindfulness-based intervention (MBI) on mental illness risks (including psychological distress, prolonged fatigue, and perceived stress) and job strain (job control and job demands) for employees with poor mental health.

Methods

A longitudinal research design was adopted. In total, 144 participants were randomized to the intervention group or the control group. The intervention group participated in MBI for eight weeks. Measurements were collected for both groups at five time points: at pre-intervention (T1), at mid-intervention (T2), at the completion of intervention (T3), four weeks after intervention (T4), and eight weeks after intervention (T5). Data were analyzed according to the intention-to-treat principle. A linear mixed model with two levels was employed to analyze the repeated measurement data.

Results

Compared with the control group, the intercepts (means at T3) for the intervention group were significantly lower on psychological distress, prolonged fatigue, and perceived stress when MBI was completed. Even with the demographic variables controlled, the positive effects remained. For growth rates of prolonged fatigue and perceived stress, participants in the intervention group showed a steeper decrease than did the participants in the control group. Regarding job strain, although the intercept (mean at T3) of job demands showed a significant decline when BMI was completed, the significance disappeared when the

demographic variables were controlled. Moreover, the other results for job control and job demands did not show promising findings.

Conclusion

As a workplace health promotion program, the MBI seems to have potential in improving mental illness risks for employees with poor mental health. However, there was insufficient evidence to support its effect on mitigating job strain. Further research on maintaining the positive effects on mental health for the long term and on developing innovative MBI to suit job strain are recommended.

Trial Registration

ClinicalTrials.gov [NCT02241070](https://clinicaltrials.gov/ct2/show/study/NCT02241070)

Introduction

A certain percentage of the world's working population has mental health-related problems. An estimated 27% of the working-age population in Europe experienced at least one type of mental health problem in the past year [1]. Mental health problems affect not only an individual's quality of life, but also his or her work capacity. Long-term mental health problems of employees may lead to individual disability and inflated expenses, such as increased absenteeism, reduced productivity, greater compensation claims and high medical costs [2–4]. Workplace health promotion is therefore needed to enhance workers' mental health. Mental health is described by WHO as “a state of well-being in which the individual realizes his or her own abilities, can cope with the normal stresses of life, can work productively and fruitfully, and is able to make a contribution to his or her community.” Due to the multifaceted definition of mental health, in order to affirm the effectiveness of a mental health promotion program, appropriate targets of assessment are necessary.

Psychological distress is common in workers, with an estimated prevalence of about 20%–30% worldwide [5–7]. In addition, prolonged fatigue and perceived stress are the most often reported symptoms among workers [8]. Prolonged fatigue is not easily reversible, not task specific, and does not disappear after a period of rest [9]. Prolonged fatigue increases vulnerability to physical and psychological problems [10]. Long-term high levels of stress are a known risk in regard to psychological distress and physiological diseases [11]. To improve workers' health, mental illness risks, such as psychological distress, prolonged fatigue, and perceived stress, need to be considered and countered.

Job strain is a notable issue in today's workforce. In Taiwan, a national survey found that more than 14% of employees “often” or “always” found their work to be very stressful [12]. The association between work stress and health problems has been well described [13,14]. The job demand-control model, based on psychosocial characteristics of work [15], is one of the most popular contemporary models for describing work stress. The model operates with two main dimensions: control at work and psychological job demands. High work demands coupled with low job control are considered to constitute job strain. However, recent research further indicates that low job control may be a better predictor of adverse health outcomes than high job demands [13]. Job control and job demands therefore could be explored separately.

Mindfulness-Based Intervention (MBI) is derived from Eastern contemplative traditions. In an operational definition, mindfulness includes two core components: (1) self-regulating attention and being curious and open, and (2) accepting one's experience at the present moment [16]. Over the past decades, previous studies regarding MBI have found impressive reductions in psychological morbidity and pain for clinical groups, and mitigation of stress and enhanced emotional well-being for nonclinical samples [16–18]. Recently, MBI has been applied to workplace mental health promotion as well. Healthcare workers were the most common subjects surveyed [19–23]. In general, these studies found that MBI was effective for psychological distress, stress, and burnout. However, most of them were carried out with a small sample size of 30–50. In addition, the results were not consistent when MBI was applied to the employees other than healthcare personnel [24–28]. Moreover, most of these previous studies collected assessment data twice or thrice, at pre- and post-intervention, which made it difficult to clarify the changes of the treatment effects in detail.

The present study therefore adopted intensive measurements at five time points to explore the effectiveness of MBI as a workplace health promotion program on mental illness risks (psychological distress, prolonged fatigue, and perceived stress) and job strain (job control and job demands) among industrial employees with a larger sample size, with and without demographic variables controlled.

Methods

The Participants and Recruitment

This study was one component of the 2011 Taiwan Workplace Mental Health Promotion Scheme. The study adopted a randomized controlled study design. Two large-scale manufacturing factories (Factory A and Factory B) were chosen for this study. All of the factories' 3270 full-time employees were requested to fill in a questionnaire comprising questions regarding mental health (measured by the Chinese Health Questionnaire, CHQ) and job strain (measured by the Job Content Questionnaire, JCQ; more details are provided in the section on Measures). Of these employees, 2849 individuals were willing to complete the questionnaire. The screening procedure was carried out between June and July in 2011. A total of 431 employees (15.13%) were found to suffer from poor mental health, which was defined by exhibiting both psychological distress and job strain in this study. That is, these employees needed to fulfill all the criteria: in the top tertile of the distribution in the CHQ for psychological distress; in the bottom tertile for the subscale of job control and in the top tertile for the subscale of job demands in the JCQ for job strain. A letter of invitation with an introduction to MBI was sent to these 431 employees. Of these, 144 responded that they would be willing to take part voluntarily in the study: 47 in Factory A and 97 in Factory B.

These 144 workers completed a baseline assessment (T1) and were then randomly allocated into the intervention group (Group I) or the waiting-list control group (Group C). To balance the sample size between groups and reduce noise or variance in the data, block randomization with a block size of four (ICIC design) was adopted for the present study [29]. In Factory A, 24 participants were allocated into Group I and 23 participants were into Group C. In Factory B, 48 participants were in Group I and 49 in Group C. In total, there were 72 participants in Group I, with the others in Group C. After randomization, two well-trained research assistants notified each participant about the group he or she was allocated to. The participants in Group C ($n = 72$) were requested to attend MBI class after completing four other times of assessment. In Group I ($n = 72$), one MBI class in Factory A ($n = 24$) and two MBI classes in Factory B ($n = 48$) were carried out to ensure an adequate class size of less than 25 for the efficiency of the intervention. All the participants were additionally measured at four other time points: at

mid-intervention (T2), at the completion of intervention (T3), four weeks after intervention (T4), and eight weeks after intervention (T5). All the measurements (T1–T5) were taken with an interval of four weeks between measurements. The research assistants distributed and collected the questionnaire. Data for this study were obtained between August and December in 2011.

Ethical Approval

This study was conducted according to the Helsinki Declaration and was approved by the Institutional Review Board of the Changhua Christian Hospital in Taiwan (IRB serial number: 110606) on July 25, 2011. Written informed consent was obtained from all the participants involved in this study. Before the Taiwanese Human Subjects Research Act was issued on Dec 28, 2011, only clinical trials according to the definition of ICH-GCP [30] were required to complete both the IRB review and clinical trial registration in Taiwan. Our study was approved by IRB and was completed prior to Dec 28, 2011. This is the reason that it was not registered prior to the enrollment of the first participant. The protocol for the trial and supporting CONSORT checklist are available as supporting information (see [S1 Protocol](#), [S2 Protocol](#) and [S1 CONSORT Checklist](#)). The authors confirm that all on-going and related trials for this intervention are registered in the ClinicalTrials.gov ([NCT02241070](#)).

Sample Size

Some open-source tools, such as GLIMMPSE, are available for calculating power and sample size for a general linear multivariate model. Mean differences in each measurement time, rather than effect size, are required [31]. However, no previous research adopting the variables compatible with our study can be referred to. In addition, multiple continuous covariates were used in this study, whereas only one continuous covariate is allowed for GLIMMPSE. GLIMMPSE therefore seems not applicable to this study. Instead, according to calculations by G*Power software, with medium effect size f equal to 0.25, type one error of .05 and 90% power being set up, a sample size of 104 participants suffices to detect a difference of means in a 2×5 mixed design ANOVA. Considering about 20–35% attrition, the 144 participants we recruited in the present research should be sufficient.

Intervention

The MBI program included two hours of mindfulness training weekly and forty-five minutes of homework practice every day for eight weeks. The participants in Group I were divided into three classes: one in Factory A and two in Factory B. The same leader and professional facilitator led the classes together in order to prevent the outcomes from being affected by the different experiences of the leader and professional facilitator. The leader was a mindfulness and vipassana meditation trainer who had practiced both types of meditation for more than two decades, and had completed mindfulness trainer education; the professional facilitator was a mindfulness practitioner and a psychiatrist. The leader was also one of the authors, to ensure the intervention provided conformed to the curriculum structure. To prevent the classes from being influenced by the leader's subjective bias, the data were kept from the leader during the period of data collection.

Each class met weekly for two hours in-session during paid working hours at the workplace. This 8-week MBI curriculum was based on the Mindfulness-Based Stress Reduction (MBSR) curriculum structure developed in the Stress Reduction Clinic of the University of Massachusetts Medical School, but without an all-day class [32]. However, the contents of the all-day class, such as silent meditation, mountain meditation, and loving-kindness meditation, were

introduced in the sessions during the eight-week course. An overview of the MBI program in this study is available as supporting information (see [S3 Protocol](#)). In general, the mindfulness training followed the template MBSR program as much as possible. The Taiwanese version of home practice CD also followed the original version developed by Kabat-Zinn [32].

A number of mindfulness techniques, including both “formal” and “informal” practices with integration in the context of Mind/Body meditation, were explored in order to cultivate moment-by-moment present awareness with non-judgmental acceptance. The awareness is also open to every single experience, regardless of whether the experience is pleasant, neutral, or unpleasant. In addition, class discussions on sharing the experiences of using the mindfulness techniques and exploring how home practices and newly acquired mindfulness skills may be used correctly were encouraged in each session. The most important aspects of the class were the leader’s and facilitator’s demonstrations of how to live mindfully every day, equipping participants to develop their own mindfulness practice independently, and identifying whether participants used problematic or maladapted strategies to face or react to challenging situations, including facing their job-related problems [33,34].

Measures

For mental illness risks. Psychological distress was measured by the Chinese Health Questionnaire (CHQ-12), a well-validated instrument [35]. The CHQ-12 was adapted from the General Health Questionnaire [36] with culturally relevant modification. The twelve items in the questionnaire, including questions pertaining to depression, anxiety, sleep disturbance, somatic concerns, and interpersonal difficulties, were on a four-point Likert scale. A higher score indicated a high level of psychological distress. For this study, alpha reliability was 0.83.

Fatigue is a common human experience. However, no gold standard or perfect definition exists for fatigue. Lewis and Wessely [37] noted several characteristics for fatigue: a subjective sensation of tiredness; having emotional, behavioral, and cognitive components; and being best viewed as a continuum. The Checklist Individual Strength questionnaire (CIS) [9] was developed to assess several aspects of prolonged fatigue: subjective fatigue, reduction in motivation, reduction in activity, and reduction in concentration; these are in line with the views of Lewis and Wessely [37]. The multidimensional CIS, which confirmed discriminant validity and convergent validity, was therefore used to measure prolonged fatigue in this study. The CIS consists of 20 items on a 7-point Likert scale [9]. The items refer to aspects of fatigue experienced during the previous 2 weeks. A higher score represented a higher level of prolonged fatigue. Cronbach’s alpha coefficient was 0.88 in the present study.

The 10-item Perceived Stress Scale (PSS-10) developed by Cohen and his colleagues was adopted to measure a global level of perceived stress [38]. The Chinese version of the PSS-10 demonstrated an adequate reliability and validity when used for a Chinese population [39]. In the present study, Cronbach’s alpha coefficient was 0.85. Participants were asked to respond to each question on a 5-point Likert scale ranging from 0 (never) to 4 (very often). A higher total score indicated a higher level of uncontrollable, unpredictable, and overwhelming feelings.

For job strain. The Job Content Questionnaire (JCQ) [40] is a commonly used questionnaire to assess job control (including 6 items pertaining to skill discretion and 3 items pertaining to decision authority), job demands (5 items), and workplace social support (8 items). The Chinese version of the JCQ [41] translated from the JCQ was reported to have acceptable psychometric properties and was adopted in this study. For the purpose of this study, only the subscales of job control and job demands were obtained. For each subscale, a sum of weighted-item scores was calculated according to the JCQ User’s Guide [40]. A higher-sum score

represented a higher level of job control or job demands. In this sample, Cronbach's alpha coefficients were 0.73 for the subscale of job control and 0.77 for the subscale of job demands.

Statistical Analysis

Data were analyzed according to the intention-to-treat principle. All participants were analyzed according to the condition (Group I or Group C) they were initially randomized. Multiple imputation by SPSS 22.0 software package was adopted for missing data. The percentages of the personal characteristics of the participants were described, and the comparisons between Group I and Group C were conducted using the chi-square test. The comparisons of unadjusted data of the dependent variables between Group I and Group C at each time point were carried out by t-test. The linear mixed model with two levels combined was employed to analyze the repeated measurement data. Level 1 was an intra-participant level and level 2 was an inter-participant level. Level 1 included only a time predictor to obtain random intercept and slope terms as outcome variables at level 2 were predicted by only intervention for Model 1, or by intervention and some demographic variables (including gender, age, education, and occupation) for Model 2. We presented the analysis results with two models for each dependent variable. In Model 1, randomly varying intercept and slope (growth rate) were explained with only the intervention variable. Due to demographic variables possibly affecting an individual's mental health, the demographic variables in level 2 were therefore controlled in Model 2. It is worthy noting that the time variable, including observations 1, 2, 3, 4, 5, was centered at median (T3) to form -2, -1, 0, 1, 2, so that the intercept term represented dependent variables at the third time measure (T3, completion of the intervention), not at the T1. The alpha level for statistical significance was set as 0.05. For easy application to workplace mental health promotion, occupation was divided into two groups: "white-collar" or "blue-collar". Women in gender and blue-collar in occupation were reference groups. All analyses were carried out using the SPSS 22.0 software package.

Results

One hundred and forty-four participants provided the initial and analyzed sample. The complete cases were made up of 58 participants in Group I (successful completion rate: 80.6%) and 54 participants in Group C (completion rate: 75.0%). The flow diagram of the study is presented in [Fig 1](#). [Table 1](#) shows the descriptive characteristics for the participants in Group I and Group C. The chi-square test revealed only statistically significant differences in gender between the groups [$\chi^2(1) = 4.85, p < .05$]. The other characteristics did not show statistical significance between the groups.

After multiple imputing of missing data, [Table 2](#) shows the comparisons of unadjusted data of mental illness risks (psychological distress, prolonged fatigue, and perceived stress) and job strain (job control and job demands) at each time point. No dependent variables at the baseline showed statistical difference between the groups. Psychological distress revealed difference at T2. At T3, all the results of mental illness risks and job strain showed significant differences between the two groups. The positive effects of MBI on mental illness risks (psychological distress, prolonged fatigue, and perceived stress) remained at T4 and at T5. In terms of job strain, after the MBI completion, only job demands showed a significant difference between the groups at T5.

The results of the linear mixed model on mental illness risks are summarized in [Table 3](#). On psychological distress, Group I had significantly lower means than Group C did at T3 in Model 1 ($\beta_{01} = -2.52, p = .001$). In Model 2, after the demographic variables were controlled, the mean difference at T3 between Group I and Group C still reached statistical significance ($\beta_{01} = -2.48$,




Fig 1. Flow diagram of MBI study.

doi:10.1371/journal.pone.0138089.g001

Table 1. Comparing the characteristics of the participants in the intervention group (Group I) and control group (Group C).

Demographic variable	Group I (n = 72)	Group C (n = 72)	χ^2 or t	p
Gender: Female, %	50.0	31.9	4.85	.028
Marital status: Married, %	73.6	70.8	0.14	.710
Education: College or above, %	83.3	80.6	0.19	.665
Occupation: White-collar, %	77.8	72.2	0.59	.441
Age, mean in year	42.4	42.7	-0.17	.863

doi:10.1371/journal.pone.0138089.t001

$p = .001$). For growth rate, the slope difference between Group I and Group C was not significant in Model 1 ($\beta_{11} = -0.54$, $p = .056$). However, in Model 2, with the demographic variables controlled, the growth rate for Group I was significantly lower than that for Group C ($\beta_{11} = -0.67$, $p = .020$).

On prolonged fatigue, Group I had significantly lower means than Group C did at T3 in both models ($\beta_{01} = -5.62$, $p = .018$ for Model 1 and $\beta_{01} = -5.59$, $p = .018$ for Model 2). The growth rates of prolonged fatigue for Group I, compared with Group C, also showed significant differences in both models ($\beta_{11} = -1.86$, $p = .016$ for Model 1 and $\beta_{11} = -2.24$, $p = .003$ for Model 2).

On perceived stress, Group I had significantly lower means than Group C did at T3 in both models ($\beta_{01} = -1.98$, $p = .005$ for Model 1 and $\beta_{01} = -1.97$, $p = .005$ for Model 2). The growth rates of perceived stress for Group I in both models also showed significant differences compared with Group C ($\beta_{11} = -0.49$, $p = .033$ for Model 1 and $\beta_{11} = -0.51$, $p = .027$ for Model 2).

The summary of the linear mixed model on job strain is shown in Table 4. On job control, no significant mean difference appeared between Group I and Group C at T3 in both models.

Table 2. Comparisons of unadjusted data of the dependent variables between Group I (n = 72) and Group C (n = 72) at each time point.

	Group	T1 [#]			T2			T3			T4			T5		
		Mean (SD)	t	p	Mean (SD)	t	p	Mean (SD)	t	p	Mean (SD)	t	p	Mean (SD)	t	p
Psychological distress			0.31	.759		-2.93	.004		-4.55	.000		-2.78	.006		-2.59	.010
	I	15.4(5.5)			10.8(6.0)			9.1(5.4)			10.3(5.6)			10.8(5.4)		
	C	15.1(5.4)			13.7(6.1)			13.7(6.7)			13.1(6.4)			13.3(6.2)		
Prolonged fatigue			-0.52	.604		-1.28	.203		-2.29	.023		-2.74	.007		-2.61	.010
	I	76.7(14.7)			73.3(15.6)			67.1(18.0)			67.5(17.8)			67.3(18.5)		
	C	78.0(13.9)			76.7(16.5)			74.2(19.2)			75.6(17.7)			75.6(19.2)		
Perceived stress			-1.60	.111		-1.08	.283		-2.85	.005		-2.92	.004		-2.83	.005
	I	19.8(4.3)			18.5(4.8)			17.3(5.4)			16.8(5.2)			16.2(5.4)		
	C	20.9(3.7)			19.4(4.7)			20.0(6.0)			19.4(5.4)			18.9(5.9)		
Job control			0.54	.592		0.05	.958		3.31	.001		1.78	.077		0.75	.456
	I	59.8(9.2)			61.8(7.9)			63.7(7.8)			62.3(9.2)			61.9(8.9)		
	C	59.0(8.3)			61.7(7.2)			59.1(8.7)			59.5(9.8)			60.7(9.8)		
Job demands			-1.00	.319		-0.11	.912		-3.27	.001		-1.89	.061		-2.15	.033
	I	32.3(5.6)			31.8(5.4)			29.0(5.3)			30.2(5.2)			30.0(5.3)		
	C	33.1(4.5)			31.9(4.8)			31.9(5.2)			32.1(6.8)			31.9(5.3)		

[#]Measured at pre-intervention (T1), at mid-intervention (T2), at the completion of intervention (T3), four weeks after intervention (T4), and eight weeks after intervention (T5).

doi:10.1371/journal.pone.0138089.t002

Table 3. Fixed effects of Intervention to the intercept and growth rate for mental illness risks.

Fixed effect	Psychological distress				Prolonged fatigue				Perceived stress			
	Model 1		Model 2		Model 1		Model 2		Model 1		Model 2	
	Coefficient (S.E.)	p	Coefficient (S.E.)	p	Coefficient (S.E.)	p	Coefficient (S.E.)	p	Coefficient (S.E.)	p	Coefficient (S.E.)	p
For intercept, π_{0i}												
Base, β_{00}	13.78(0.53)	< .001	13.80(1.24)	< .001	76.00(1.65)	< .001	78.85(3.81)	< .001	19.71(0.49)	< .001	19.87(1.13)	< .001
Intervention, β_{01}	-2.52(0.74)	.001	-2.48(0.76)	.001	-5.62(2.34)	.018	-5.59(2.34)	.018	-1.98(0.69)	.005	-1.97(0.70)	.005
Gender, β_{02}			0.27(0.87)	.757			-0.65(2.68)	.808			0.19(0.80)	.813
Age, β_{03}			0.01(0.05)	.815			-0.37(0.15)	.013			-0.10(0.04)	.018
Education, β_{04}			-0.65(0.78)	.406			1.15(2.39)	.633			-0.39(0.71)	.589
Occupation, β_{05}			-0.33(1.01)	.742			-3.17(3.10)	.309			-0.41(0.92)	.659
For growth rate, π_{1i}												
Time, β_{10}	-0.43(0.20)	.030	0.36(0.46)	.430	-0.59(0.54)	.281	2.69(1.21)	.028	-0.41(0.16)	.012	0.18(0.37)	.639
Intervention×Time, β_{11}	-0.54(0.28)	.056	-0.67(0.28)	.020	-1.86(0.77)	.016	-2.24(0.75)	.003	-0.49(0.23)	.033	-0.51(0.23)	.027
Gender×Time, β_{12}			-0.86(0.32)	.008			-3.16(0.85)	< .001			-0.41(0.26)	.121
Age×Time, β_{13}			0.02(0.02)	.198			0.11(0.05)	.017			0.02(0.01)	.216
Education×Time, β_{14}			0.14(0.29)	.624			1.56(0.76)	.043			0.42(0.23)	.076
Occupation×Time, β_{15}			-0.28(0.37)	.449			-1.46(0.99)	.142			-0.39(0.30)	.201

Note: Time variable was recoded as -2, -1, 0, 1, 2, for five measurement time points in order. Women in Gender, blue collar in Occupation, and control group in Intervention were reference groups. Age was centered at mean and Education was centered at the level of college. The level 1 included a time variable only as the predictor, and level 2 included Intervention, Gender, Age, Education, and Occupation as predictors. For simplicity, variance components of level 1 and level 2 were omitted in the Table.

Each model was combined with level 1 and level 2 equations. For level 1 equation: $Y_{ti} = \pi_{0i} + \pi_{1i}Time_{ti} + e_{ti}$. For level 2 equation:

$\pi_{0i} = \beta_{00} + \beta_{01}Intervention_i + r_{0i}$ and $\pi_{1i} = \beta_{10} + \beta_{11}Intervention_i + r_{1i}$ in Model 1;

$\pi_{0i} = \beta_{00} + \beta_{01}Intervention_i + \beta_{02}Gender_i + \beta_{03}Age_i + \beta_{04}Education_i + \beta_{05}Occupation_i + r_{0i}$ and

$\pi_{1i} = \beta_{10} + \beta_{11}Intervention_i + \beta_{12}Gender_i + \beta_{13}Age_i + \beta_{14}Education_i + \beta_{15}Occupation_i + r_{1i}$ in Model 2.

doi:10.1371/journal.pone.0138089.t003

The growth rates of job control for Group I, compared with Group C, also showed no significant difference in both models.

Regarding job demands, Group I had significantly lower means than Group C did at T3 in Model 1 ($\beta_{01} = -1.52, p = .043$); however, no significant difference appeared for the growth rate between Group I and Group C. In Model 2 with the demographic variables controlled, the mean difference at T3 disappeared ($\beta_{01} = -1.50, p = .054$). The growth rate of job demands in Model 2 did not show significant difference between the two groups as well.

Discussion

The present study adopted a randomized controlled study design to determine the effectiveness of MBI as a workplace health promotion program. Compared with the control group, the intervention effects on psychological distress, prolonged fatigue, and perceived stress were revealed when the intervention was completed (T3). Furthermore, the positive effects on prolonged fatigue and perceived stress were maintained with time. However, in general, the effects of MBI on job control and job demands were not statistically significant. These findings show that

Table 4. Fixed effects of Intervention to the intercept and growth rate for job strain.

Fixed effect	Job control				Job demands			
	Model 1		Model 2		Model 1		Model 2	
	Coefficient (S.E.)	p	Coefficient (S.E.)	p	Coefficient (S.E.)	p	Coefficient (S.E.)	p
For intercept, π_{0i}								
Base, β_{00}	60.00(0.89)	< .001	58.60(2.04)	< .001	32.17(0.53)	< .001	32.10(1.25)	< .001
Intervention, β_{01}	1.87(1.25)	.137	1.92(1.26)	.128	-1.52(0.75)	.043	-1.50(0.77)	.054
Gender, β_{02}			0.40(1.43)	.782			0.07(0.88)	.936
Age, β_{03}			0.21(0.08)	.009			0.00(0.05)	.919
Education, β_{04}			0.14(1.28)	.913			0.14(0.79)	.855
Occupation, β_{05}			1.53(1.66)	.358			0.04(1.02)	.970
For growth rate, π_{1i}								
Time, β_{10}	0.11(0.25)	.660	-1.68(0.55)	.003	-0.24(0.16)	.149	-0.01(0.38)	.980
Intervention \times Time, β_{11}	0.35(0.35)	.315	0.53(0.34)	.116	-0.39(0.23)	.093	-0.41(0.24)	.086
Gender \times Time, β_{12}			1.55(0.39)	< .001			-0.22(0.27)	.406
Age \times Time, β_{13}			-0.05(0.02)	.012			0.02(0.01)	.119
Education \times Time, β_{14}			-0.53(0.35)	.130			0.22(0.24)	.370
Occupation \times Time, β_{15}			0.99(0.45)	.029			-0.09(0.31)	.775

Note: Time variable was recoded as -2, -1, 0, 1, 2, for five measurement time points in order. Women in Gender, blue collar in Occupation, and control group in Intervention were reference groups. Age was centered at mean and Education was centered at the level of college. The level 1 included a time variable only as the predictor, and level 2 included Intervention, Gender, Age, Education, and Occupation as predictors. For simplicity, variance components of level 1 and level 2 were omitted in the Table.

Each model was combined with level 1 and level 2 equations. For level 1 equation: $Y_{ti} = \pi_{0i} + \pi_{1i}Time_{ti} + e_{ti}$. For level 2 equation:

$\pi_{0i} = \beta_{00} + \beta_{01}Intervention_i + r_{0i}$ and $\pi_{1i} = \beta_{10} + \beta_{11}Intervention_i + r_{1i}$ in Model 1;

$\pi_{0i} = \beta_{00} + \beta_{01}Intervention_i + \beta_{02}Gender_i + \beta_{03}Age_i + \beta_{04}Education_i + \beta_{05}Occupation_i + r_{0i}$ and

$\pi_{1i} = \beta_{10} + \beta_{11}Intervention_i + \beta_{12}Gender_i + \beta_{13}Age_i + \beta_{14}Education_i + \beta_{15}Occupation_i + r_{1i}$ in Model 2.

doi:10.1371/journal.pone.0138089.t004

MBI has the potential to lessen the mental illness risks for the workers with poor mental health, but not for job strain.

The literature includes limited reports on the use of MBI to improve workers' mental health. Most of the reports showed results similar to ours. Studies using MBI for healthcare personnel showed that MBI was an effective strategy not only for reducing burnout and perceived stress, but also for increasing patients' satisfaction [19–23]. The qualitative findings also revealed that greater relaxation and patience, more self-care and self-acceptance, improved inner peace, compassion and joy, better focus, and fewer somatic symptoms were the reported benefits [42,43]. In addition, research using a shortened worksite MBI showed improvement in perceived stress, psychological distress, positive orientation to life, and sleep quality among working adults [20,23,44,45]. How may MBI enhance mental health? Through the practice of meditation, individuals can achieve a state of mindfulness, in which thoughts and feelings are observed as events in the mind; any negative feeling or thought then become decentralized as “an event.” This dispassionate state of self-observation might create a “space” between one's perception and response, helping to enhance emotional well-being and mental health [17].

However, not all the studies agree with our positive results. A worksite mindfulness-related multi-component health promotion intervention found no effect on work engagement, mental health, need for recovery, and mindfulness after 6 and 12 months of intervention [24]. Some possibilities might explain the inconsistency between this and our findings. First, in the study done by van Berkel et al. [24], all the voluntary participants were enrolled, so the ceiling effect

might be the reason why the dependent variables could not be improved. Evidence was found of psychologically healthy participants diluting the observed effects of worksite stress management training programs [46]. An effective workplace health-promoting program needs to be tailored to the workers' physical, psychological, or social needs [47]. Our study thus targeted the participants with poor mental health, and MBI was adopted to enhance their mental health. Second, lack of effects in the study of van Berkel et al. [24] might be due to the intervention's being run on the participants' own time. Our intervention was implemented during paid working hours. In other studies with positive results, their MBIs were also delivered at the workplace during work hours [20,22,25]. Effective workplace health-promoting interventions need the organization's commitment to facilitating the changes both for employees and their working conditions [47]. This is the same as the action strategy of creating supportive environments as set forth in the Ottawa Charter of Health Promotion [48]. Another reason may be the different follow-up time. Based on the results of our study, the immediate and short-term positive effects of MBI on mental illness risks were observed. Determining how to modify the program and extend the positive effect to a longer period would be a challenge in the future.

Job control and job demands did not show significant improvements with time in Group I. The MBI class took place in working hours, and job content did not change, which might create other demanding characteristics. It might be one of the reasons why MBI did not work out in this field. However, there is a paucity of research on the relations between the level of mindfulness and either job control or job demands. Actual explanations are still unknown. Previous studies using MBI to improve work-related factors focused mainly on burnout and job satisfaction. The findings generally showed beneficial results in decreasing burnout and enhancing job satisfaction [20,22,49,50]. However, when MBI was adopted as an intervention for the outcomes which are not emotion-related, such as work engagement, lifestyle behaviors, and blood inflammation markers; all the results did not show significant differences [24,26,27]. The traditional MBI seemed to take effect mainly based on emotions or feelings. Besides, previous studies which tried MBI as an auxiliary strategy in eating behaviors and smoking cessation achieved some favorable results [51,52]. In the future, MBI with some innovations to apply to work-related dimensions or to health behaviors are recommended.

Notably, there were significant effects of time on psychological distress, prolonged fatigue, and perceived stress, with or without demographic variables controlled. It implies that both groups improved on mental illness risks with time; the participants in Group C also reduced psychological distress, fatigue, and stress, although the participants in Group I decreased more sharply. A possible explanation for this is that the participants in Group C also felt being cared for by being invited to take part in assessment and completing the questionnaire regarding mental health every four weeks. It implies that other simple programs to enhance workers' mental health are worth developing in addition to MBI. In addition, the demographic variables showed their own respective effects on dependent variables. Decades of evidence have consistently shown that mental health is not equally distributed across socio-economic strata or gender [53–55]. However, previous studies using MBI as an intervention in workplace health promotion did not consider the influences of these variables. The present study attests to the significance of demographic variables, such as gender, age, and educational level, on mental health when implementing MBI for workplace health promotion.

The major strength of this study is the use of a randomized controlled study design with intensive repeated measures. This reliable research design could show the changes of dependent variables in detail with time. Second, the measured variables were multifaceted, matching the variety of mental health facets. Third, the sample size of the intervention group was relatively large compared with previous randomized controlled trials that generally included approximately twenty participants in the intervention. These aspects may avoid some research

errors. However, some limitations in the study arose as well. First, most of the participants were married, highly educated, and white-collar. The inference of the findings should be applied with caution. Second, the extent of home meditation practice was thought to be associated with the effects of the MBI [56]. The study did not quantify and analyze the extent of home practice. Third, the waiting-list control group was not active. Some nonspecific factors, such as therapists' attention, social support, and positive expectancy, cannot be excluded. The improved outcomes may be partly attributed to these factors. In addition, the study was not conducted in a blind study design; such study design did not prevent research outcomes from being influenced by the expectations of the participants in Group I. Finally, although one of the authors being the leader of MBI could ensure that the study was conducted smoothly and the intervention conformed to the curriculum structure, such design may entail the possibility of a conflict of interest and cannot eliminate subjective bias on the part of the leader. Further studies adopting active control groups with a double-blind design are thus recommended.

Supporting Information

S1 CONSORT Checklist. CONSORT Checklist.
(DOC)

S1 Dataset. The dataset of this study.
(XLS)

S1 Protocol. Trial Protocol in Chinese.
(DOC)

S2 Protocol. Trial Protocol in English.
(DOC)

S3 Protocol. MBI program protocol.
(DOC)

Acknowledgments

We are grateful for the cooperation of the factories, and we thank the workers who participated in the study and provided valuable information.

Author Contributions

Conceived and designed the experiments: SLH FYH FCT. Performed the experiments: SLH FYH FCT. Analyzed the data: RHL SLH. Contributed reagents/materials/analysis tools: RHL FCT SLH. Wrote the paper: SLH RHL FYH FCT.

References

1. WHO Regional Office for Europe. Mental health: Data and statistics. 2014. Available: <http://www.euro.who.int/en/health-topics/noncommunicable-diseases/mental-health/data-and-statistics>.
2. International Labour Office. Mental health in the workplace. 2000. Available: http://www.ilo.org/wcmsp5/groups/public/-ed_emp/-ifp_skills/documents/publication/wcms_108221.pdf.
3. Siegrist J. Stress at work. In: Smelser NJ, Baltes PB, editors. International encyclopedia of the social & behavioral sciences. New York: Elsevier Ltd; 2001. pp. 15175–15179.
4. Goetzel RZ, Pei X, Tabrizi MJ, Henke RM, Kowlessar N, Nelson CF, et al. Ten modifiable health risk factors are linked to more than one-fifth of employer-employee health care spending. *Health Aff.* 2012; 31: 2474–2484.

5. Goodwin L, Ben-Zion I, Fear NT, Hotopf M, Stansfeld SA, Wessely S. Are reports of psychological stress higher in occupational studies? A systematic review across occupational and population based studies. *PLoS One*. 2013; 8: e78693. doi: [10.1371/journal.pone.0078693](https://doi.org/10.1371/journal.pone.0078693) PMID: [24223840](https://pubmed.ncbi.nlm.nih.gov/24223840/)
6. Marchand A, Demers A, Durand P. Do occupation and work conditions really matter? A longitudinal analysis of psychological distress experiences among Canadian workers. *Social Health Illn*. 2005; 27: 602–627. PMID: [16078903](https://pubmed.ncbi.nlm.nih.gov/16078903/)
7. Huang SL, Lee HS, Li RH, Lai YM, Chen AL, Tang FC. Differences in health complaints among Taiwanese workers in different occupational categories. *J Occup Health*. 2012; 54: 241–249. PMID: [22790527](https://pubmed.ncbi.nlm.nih.gov/22790527/)
8. Agnès PT, Enrique FM, John H, Greet V. Fourth European working conditions survey (2005). 2010. Available: <http://www.eurofound.europa.eu/ewco/surveys/ewcs2005/index.htm>.
9. Beurskens AJ, Bultmann U, Kant I, Vercoulen JH, Bleijenberg G, Swaen GM. Fatigue among working people: validity of a questionnaire measure. *Occup Environ Med*. 2000; 57: 353–357. PMID: [10769302](https://pubmed.ncbi.nlm.nih.gov/10769302/)
10. Leone SS, Huibers MJ, Knottnerus JA, Kant I. Similarities, overlap and differences between burnout and prolonged fatigue in the working population. *QJM*. 2007; 100: 617–627. PMID: [17921196](https://pubmed.ncbi.nlm.nih.gov/17921196/)
11. Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *Jama*. 2007; 298: 1685–1687. PMID: [17925521](https://pubmed.ncbi.nlm.nih.gov/17925521/)
12. Hsu JH, Wu YS. Survey of perceptions of safety and health in the work environment in 2010 Taiwan. Taipei: Institute of Labor, Occupational Safety and Health, Ministry of Labor; 2011.
13. Meyer JD, Cifuentes M, Warren N. Association of self-rated physical health and incident hypertension with O NET factors: validation using a representative national survey. *J Occup Environ Med*. 2011; 53: 139–145. PMID: [21270664](https://pubmed.ncbi.nlm.nih.gov/21270664/)
14. Meyer JD. Race-based job discrimination, disparities in job control, and their joint effects on health. *Am J Ind Med*. 2014; 57: 587–595. doi: [10.1002/ajim.22255](https://doi.org/10.1002/ajim.22255) PMID: [24105870](https://pubmed.ncbi.nlm.nih.gov/24105870/)
15. Karasek RA Jr. Job demands, job decision latitude, and mental strain: Implications for job redesign. *Adm Sci Q*. 1979: 285–308.
16. Bishop SR, Lau M, Shapiro S, Carlson L, Anderson ND, Carmody J, et al. Mindfulness: A proposed operational definition. *Clinical Psychology: Science and Practice*. 2004; 11: 230–241.
17. Grossman P, Niemann L, Schmidt S, Walach H. Mindfulness-based stress reduction and health benefits. A meta-analysis. *J Psychosom Res*. 2004; 57: 35–43. PMID: [15256293](https://pubmed.ncbi.nlm.nih.gov/15256293/)
18. Lakhan SE, Schofield KL. Mindfulness-based therapies in the treatment of somatization disorders: a systematic review and meta-analysis. *PLoS One*. 2013; 8: e71834. doi: [10.1371/journal.pone.0071834](https://doi.org/10.1371/journal.pone.0071834) PMID: [23990997](https://pubmed.ncbi.nlm.nih.gov/23990997/)
19. Cohen-Katz J, Wiley SD, Capuano T, Baker DM, Kimmel S, Shapiro S. The effects of mindfulness-based stress reduction on nurse stress and burnout, Part II: A quantitative and qualitative study. *Holist Nurs Pract*. 2005; 19: 26–35. PMID: [15736727](https://pubmed.ncbi.nlm.nih.gov/15736727/)
20. Duchemin AM, Steinberg BA, Marks DR, Vanover K, Klatt M. A small randomized pilot study of a workplace mindfulness-based intervention for surgical intensive care unit personnel: effects on salivary alpha-amylase levels. *J Occup Environ Med*. 2015; 57: 393–399. PMID: [25629803](https://pubmed.ncbi.nlm.nih.gov/25629803/)
21. Bazarko D, Cate RA, Azocar F, Kreitzer MJ. The Impact of an Innovative Mindfulness-Based Stress Reduction Program on the Health and Well-Being of Nurses Employed in a Corporate Setting. *J Workplace Behav Health*. 2013; 28: 107–133. PMID: [23667348](https://pubmed.ncbi.nlm.nih.gov/23667348/)
22. Horner JK, Piercy BS, Eure L, Woodard EK. A pilot study to evaluate mindfulness as a strategy to improve inpatient nurse and patient experiences. *Appl Nurs Res*. 2014; 27: 198–201. doi: [10.1016/j.apnr.2014.01.003](https://doi.org/10.1016/j.apnr.2014.01.003) PMID: [24602399](https://pubmed.ncbi.nlm.nih.gov/24602399/)
23. Pipe TB, Bortz JJ, Dueck A, Pendergast D, Buchda V, Summers J. Nurse leader mindfulness meditation program for stress management: a randomized controlled trial. *J Nurs Adm*. 2009; 39: 130–137. PMID: [19590469](https://pubmed.ncbi.nlm.nih.gov/19590469/)
24. van Berkel J, Boot CR, Proper KI, Bongers PM, van der Beek AJ. Effectiveness of a worksite mindfulness-related multi-component health promotion intervention on work engagement and mental health: results of a randomized controlled trial. *PLoS One*. 2014; 9: e84118. doi: [10.1371/journal.pone.0084118](https://doi.org/10.1371/journal.pone.0084118) PMID: [24489648](https://pubmed.ncbi.nlm.nih.gov/24489648/)
25. Aikens KA, Astin J, Pelletier KR, Levanovich K, Baase CM, Park YY, et al. Mindfulness goes to work: impact of an online workplace intervention. *J Occup Environ Med*. 2014; 56: 721–731. PMID: [24988100](https://pubmed.ncbi.nlm.nih.gov/24988100/)
26. van Berkel J, Boot CR, Proper KI, Bongers PM, van der Beek AJ. Effectiveness of a worksite mindfulness-based multi-component intervention on lifestyle behaviors. *The international journal of behavioral nutrition and physical activity*. 2014; 11: 9. doi: [10.1186/1479-5868-11-9](https://doi.org/10.1186/1479-5868-11-9) PMID: [24467802](https://pubmed.ncbi.nlm.nih.gov/24467802/)

27. Malarkey WB, Jarjoura D, Klatt M. Workplace based mindfulness practice and inflammation: a randomized trial. *Brain Behav Immun*. 2013; 27: 145–154. doi: [10.1016/j.bbi.2012.10.009](https://doi.org/10.1016/j.bbi.2012.10.009) PMID: [23078984](https://pubmed.ncbi.nlm.nih.gov/23078984/)
28. Wolever RQ, Bobinet KJ, McCabe K, Mackenzie ER, Fekete E, Kusnick CA, et al. Effective and viable mind-body stress reduction in the workplace: a randomized controlled trial. *J Occup Health Psychol*. 2012; 17: 246–258. doi: [10.1037/a0027278](https://doi.org/10.1037/a0027278) PMID: [22352291](https://pubmed.ncbi.nlm.nih.gov/22352291/)
29. Lachin JM, Matts JP, Wei L. Randomization in clinical trials: conclusions and recommendations. *Control Clin Trials*. 1988; 9: 365–374. PMID: [3203526](https://pubmed.ncbi.nlm.nih.gov/3203526/)
30. ICH GCP. Glossary. 2014. Available: <http://ichgcp.net/1-glossary>.
31. Guo Y, Logan HL, Glueck DH, Muller KE. Selecting a sample size for studies with repeated measures. *BMC Med Res Methodol*. 2013; 13: 100. doi: [10.1186/1471-2288-13-100](https://doi.org/10.1186/1471-2288-13-100) PMID: [23902644](https://pubmed.ncbi.nlm.nih.gov/23902644/)
32. Kabat-Zinn J. Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness. New York: Delta; 1990.
33. Dobkin PL, Hickman S, Monshat K. Holding the heart of Mindfulness-Based Stress Reduction: Balancing fidelity and imagination when adapting MBSR. *Mindfulness*. 2014; 5: 710–718.
34. Blacker M, Meleo-Meyer F, Kabat-Zinn J, Santorelli S. Stress reduction clinic Mindfulness-Based Stress Reduction (MBSR) curriculum guide. Worcester: University of Massachusetts Medical School; 2009.
35. Cheng TA, Williams P. The design and development of a screening questionnaire (CHQ) for use in community studies of mental disorders in Taiwan. *Psychol Med*. 1986; 16: 415–422. PMID: [3726013](https://pubmed.ncbi.nlm.nih.gov/3726013/)
36. Goldberg DP. Manual of the general health questionnaire. Windsor: NFER; 1978.
37. Lewis G, Wessely S. The epidemiology of fatigue: more questions than answers. *J Epidemiol Community Health*. 1992; 46: 92–97. PMID: [1583440](https://pubmed.ncbi.nlm.nih.gov/1583440/)
38. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav*. 1983; 24: 385–396. PMID: [6668417](https://pubmed.ncbi.nlm.nih.gov/6668417/)
39. Wang Z, Chen J, Boyd JE, Zhang H, Jia X, Qiu J, et al. Psychometric properties of the Chinese version of the Perceived Stress Scale in policewomen. *PLoS One*. 2011; 6: e28610. doi: [10.1371/journal.pone.0028610](https://doi.org/10.1371/journal.pone.0028610) PMID: [22164311](https://pubmed.ncbi.nlm.nih.gov/22164311/)
40. Karasek RA. Job content questionnaire and user's guide. Lowell: University of Massachusetts; 1985.
41. Cheng Y, Luh WM, Guo YL. Reliability and validity of the Chinese version of the Job Content Questionnaire in Taiwanese workers. *Int J Behav Med*. 2003; 10: 15–30. PMID: [12581945](https://pubmed.ncbi.nlm.nih.gov/12581945/)
42. Cohen-Katz J, Wiley S, Capuano T, Baker DM, Deitrick L, Shapiro S. The effects of mindfulness-based stress reduction on nurse stress and burnout: a qualitative and quantitative study, part III. *Holist Nurs Pract*. 2005; 19: 78–86. PMID: [15871591](https://pubmed.ncbi.nlm.nih.gov/15871591/)
43. Moody K, Kramer D, Santizo RO, Magro L, Wyshogrod D, Ambrosio J, et al. Helping the helpers: mindfulness training for burnout in pediatric oncology—a pilot program. *J Pediatr Oncol Nurs*. 2013; 30: 275–284. doi: [10.1177/1043454213504497](https://doi.org/10.1177/1043454213504497) PMID: [24101747](https://pubmed.ncbi.nlm.nih.gov/24101747/)
44. Klatt MD, Buckworth J, Malarkey WB. Effects of low-dose mindfulness-based stress reduction (MBSR-I) on working adults. *Health Educ Behav*. 2009; 36: 601–614. doi: [10.1177/1090198108317627](https://doi.org/10.1177/1090198108317627) PMID: [18469160](https://pubmed.ncbi.nlm.nih.gov/18469160/)
45. Foureur M, Besley K, Burton G, Yu N, Crisp J. Enhancing the resilience of nurses and midwives: pilot of a mindfulness-based program for increased health, sense of coherence and decreased depression, anxiety and stress. *Contemp Nurse*. 2013; 45: 114–125. doi: [10.5172/conu.2013.45.1.114](https://doi.org/10.5172/conu.2013.45.1.114) PMID: [24099232](https://pubmed.ncbi.nlm.nih.gov/24099232/)
46. Flaxman PE, Bond FW. Worksite stress management training: moderated effects and clinical significance. *J Occup Health Psychol*. 2010; 15: 347–358. doi: [10.1037/a0020522](https://doi.org/10.1037/a0020522) PMID: [21058850](https://pubmed.ncbi.nlm.nih.gov/21058850/)
47. Czabala C, Charzynska K, Mroziak B. Psychosocial interventions in workplace mental health promotion: an overview. *Health Prom Int*. 2011; 26: 70–84.
48. WHO. Ottawa charter for health promotion. 1986. Available: <http://www.who.int/healthpromotion/conferences/previous/ottawa/en/>.
49. Hülshager UR, Alberts HJ, Feinholdt A, Lang JW. Benefits of mindfulness at work: The role of mindfulness in emotion regulation, emotional exhaustion, and job satisfaction. *J Appl Psychol*. 2013; 98: 310. doi: [10.1037/a0031313](https://doi.org/10.1037/a0031313) PMID: [23276118](https://pubmed.ncbi.nlm.nih.gov/23276118/)
50. Fortney L, Luchterhand C, Zakletskaia L, Zgierska A, Rakei D. Abbreviated mindfulness intervention for job satisfaction, quality of life, and compassion in primary care clinicians: a pilot study. *Ann Fam Med*. 2013; 11: 412–420. doi: [10.1370/afm.1511](https://doi.org/10.1370/afm.1511) PMID: [24019272](https://pubmed.ncbi.nlm.nih.gov/24019272/)
51. Bush HE, Rossy L, Mintz LB, Schopp L. Eat for life: a work site feasibility study of a novel mindfulness-based intuitive eating intervention. *Am J Health Promot*. 2014; 28: 380–388. doi: [10.4278/ajhp.120404-QUAN-186](https://doi.org/10.4278/ajhp.120404-QUAN-186) PMID: [23941103](https://pubmed.ncbi.nlm.nih.gov/23941103/)

52. Michalsen A, Richarz B, Reichardt H, Spahn G, Konietzko N, Dobos GJ. [Smoking cessation for hospital staff. A controlled intervention study]. *Dtsch Med Wochenschr.* 2002; 127: 1742–1747. PMID: [12192632](#)
53. Dohrenwend BP. Socioeconomic status (SES) and psychiatric disorders. Are the issues still compelling? *Soc Psychiatry Psychiatr Epidemiol.* 1990; 25: 41–47. PMID: [2406949](#)
54. Wang PS, Berglund P, Kessler RC. Recent care of common mental disorders in the United States: prevalence and conformance with evidence-based recommendations. *J Gen Intern Med.* 2000; 15: 284–292. PMID: [10840263](#)
55. Afifi M. Gender differences in mental health. *Singapore Med J.* 2007; 48: 385–391. PMID: [17453094](#)
56. Rosenzweig S, Greeson JM, Reibel DK, Green JS, Jasser SA, Beasley D. Mindfulness-based stress reduction for chronic pain conditions: variation in treatment outcomes and role of home meditation practice. *J Psychosom Res.* 2010; 68: 29–36. doi: [10.1016/j.jpsychores.2009.03.010](#) PMID: [20004298](#)